

## OriGene Technologies Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850

Phone: +1-888-267-4436 Fax: +1-301-340-8606 techsupport@origene.com

**UNITED STATES** 

SM2091RT

OriGene EU

## official Lo

**Acris Antibodies GmbH** 

Schillerstr. 5 32052 Herford GERMANY

Phone: +49-5221-34606-0 Fax: +49-5221-34606-11 info@acris-antibodies.com

## Monoclonal Antibody to NKG2A/C/E - PE

Alternate names: CD159a, CD159c, CD159e, NKG2 lectin-like family

Catalog No.: SM2091RT
Quantity: 25 Tests
Concentration: 0.1 mg/ml

Background: In mice, NKG2 subunits associate with CD94 to form heterodimers at the surface of natural

killer (NK) cells. The CD94/NKG2 heterodimer is the receptor for a non-classical MHC class I

ligand, which is Qa-1 in the mouse.

Host / Isotype: Rat / IgG2a Clone: 20d5

Immunogen: CHO transfected cells expressing the B6 allele of NKG2A.

Spleen cells from immunised Lewis rats were fused with cells of the mouse P3X63-Ag8.653

myeloma cell line.

Format: State: Lyophilized purified IgG fraction.

Purification: Affinity Chromatography on Protein G.

**Buffer System:** PBS, pH7.4 containing 0.09% Sodium Azide and 1% Bovine Serum Albumin.

**Label:** PE – R. Phycoerythrin (RPE)

**Reconstitution:** Restore with 1 ml distilled water

Applications: Flow Cytometry: Neat - 1/2; Use 10µl of the suggested working dilution to label 10e6 cells in

100µl; The Fc region of monoclonal antibodies may bind non-specifically to cells

expressing low affinity fc receptors.

Other applications not tested. Optimal dilutions are dependent on conditions and should

be determined by the user.

Specificity: This antibody recognises NKG2A, NKG2C and NKG2E, which are isoforms of the NKG2 lectin-

like family.

Clone 20d5 is reported to block ligand binding to the receptor. We recommend the use of

SM2091LE for this purpose.

Species: Mouse.

Other species not tested.

**Storage:** Store the antibody undiluted at 2-8°C.

DO NOT FREEZE!

This product is photosensitive and should be protected from light.

Shelf life: one year from despatch.

General Readings: 1. Vance RE, Jamieson AM, Raulet DH. Recognition of the class Ib molecule Qa-1(b) by

putative activating receptors CD94/NKG2C and CD94/NKG2E on mouse natural killer cells. J

Exp Med. 1999 Dec 20;190(12):1801-12. PubMed PMID: 10601355.

2. Vance RE, Jamieson AM, Cado D, Raulet DH. Implications of CD94 deficiency and

For research and in vitro use only. Not for diagnostic or therapeutic work.

Material Safety Datasheets are available at www.acris-antibodies.com or on request.

TUV NORD
TUV NORD CERT
GribH EN ISO 965



## SM2091RT: Monoclonal Antibody to NKG2A/C/E - PE

monoallelic NKG2A expression for natural killer cell development and repertoire formation. Proc Natl Acad Sci U S A. 2002 Jan 22;99(2):868-73. Epub 2002 Jan 8. PubMed PMID: 11782535.

