

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850 UNITED STATES Phone: +1-888-267-4436 Fax: +1-301-340-8606 techsupport@origene.com OriGene Technologies GmbH

Schillerstr. 5 32052 Herford GERMANY Phone: +49-5221-34606-0 Fax: +49-5221-34606-11 info-de@origene.com

SM2093R Monoclonal Antibody to CD16/CD32 - PE


Alternate names:	CD16a, FCG3, FCGR3A, Fc gamma receptor IIB, Fc-gamma RIII, Fc-gamma RIII-alpha, Fc-gamma RIIIa, Fc-gamma-RIIB, FcR mouse Seroblock, FcRII, FcRIII, FcRIIIa, Fcgr2, IGFR3, IgG Fc receptor III-2, Low affinity immunoglobulin gamma Fc region receptor III-A, Low affinity immunoglobulin gamma Fc region receptor III-B, Ly-17
Quantity:	100 Tests
Concentration:	0.1 mg/ml
Background:	In the mouse only the transmembrane form of CD16 is reported to exist, which is expressed on macrophages, NK cells, granulocytes, myeloid precursors, and subpopulations of T lymphocytes. CD32 is primarily expressed on cells of the myeloid lineage and also on mature B lymphocytes.
Host / Isotype:	Rat / IgG2b
Recommended Isotype Controls:	SM19R
Clone:	FCR 4G8
Immunogen:	PU5 1.8 IOE7 Balb/c mouse cell line. Cells from immunised rats were fused with cells of the mouse P3X63-Ag8.653 myeloma cell line.
Format:	 State: Lyophilized purified IgG Purification: Affinity chromatography on Protein G Buffer System: PBS, pH7.4 containing 0.09% Sodium Azide and 1% Bovine Serum Albumin Label: PE – R. Phycoerythrin (RPE) Reconstitution: Restore with 1 ml distilled water
Applications:	Flow Cytometry: Neat - 1/10; Use 10µl of the suggested working dilution to label 10e6 cells in 100µl. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Specificity:	This antibody recognises an epitope expressed by the low affinity Fc receptors, Fc gamma III (CD16) and Fc gamma II (CD32). Species: Mouse. Other species not tested.
Storage:	Store the antibody undiluted at 2-8°C. DO NOT FREEZE! This product is photosensitive and should be protected from light. Shelf life: one year from despatch.

For research and in vitro use only. Not for diagnostic or therapeutic work. Material Safety Datasheets are available at www.acris-antibodies.com or on request.



Pictures:

Staining of mouse peritoneal macrophages with RAT ANTI MOUSE CD16/32 : RPE (SM2093R).



For research and in vitro use only. Not for diagnostic or therapeutic work. Material Safety Datasheets are available at www.acris-antibodies.com or on request.