

**SM2093RT****Monoclonal Antibody to CD16/CD32 - PE**

<b>Alternate names:</b>	CD16a, FCG3, FCGR3A, Fc gamma receptor IIB, Fc-gamma RIII, Fc-gamma RIII-alpha, Fc-gamma RIIIa, Fc-gamma-RIIB, FcR mouse Seroblock, FcRII, FcRIII, FcRIIIa, Fcgr2, IGFR3, IgG Fc receptor III-2, Low affinity immunoglobulin gamma Fc region receptor III-A, Low affinity immunoglobulin gamma Fc region receptor III-B, Ly-17
<b>Quantity:</b>	25 Tests
<b>Concentration:</b>	0.1 mg/ml
<b>Background:</b>	In the mouse only the transmembrane form of CD16 is reported to exist, which is expressed on macrophages, NK cells, granulocytes, myeloid precursors, and subpopulations of T lymphocytes. CD32 is primarily expressed on cells of the myeloid lineage and also on mature B lymphocytes.
<b>Host / Isotype:</b>	Rat / IgG2b
<b>Recommended Isotype Controls:</b>	SM19R
<b>Clone:</b>	FCR4G8
<b>Immunogen:</b>	PU5 1.8 IOE7 Balb/c mouse cell line. Cells from immunised rats were fused with cells of the mouse P3X63-Ag8.653 myeloma cell line.
<b>Format:</b>	<b>State:</b> Lyophilized purified IgG <b>Purification:</b> Affinity chromatography on Protein G <b>Buffer System:</b> PBS, pH7.4 containing 0.09% Sodium Azide and 1% Bovine Serum Albumin <b>Label:</b> PE – R. Phycoerythrin (RPE) <b>Reconstitution:</b> Restore with 1 ml distilled water
<b>Applications:</b>	Flow Cytometry: Neat - 1/10; Use 10µl of the suggested working dilution to label 10e6 cells in 100µl. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
<b>Specificity:</b>	This antibody recognises an epitope expressed by the low affinity Fc receptors, Fc gamma III (CD16) and Fc gamma II (CD32). <b>Species:</b> Mouse. Other species not tested.
<b>Storage:</b>	Store the antibody undiluted at 2-8°C. DO NOT FREEZE! This product is photosensitive and should be protected from light. Shelf life: one year from despatch.

## Pictures:

Staining of mouse peritoneal macrophages with RAT ANTI MOUSE CD16/32 : RPE (SM2093R).

