

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850 UNITED STATES Phone: +1-888-267-4436 Fax: +1-301-340-8606 techsupport@origene.com

OriGene Technologies GmbH

Schillerstr. 5 32052 Herford GERMANY Phone: +49-5221-34606-0 Fax: +49-5221-34606-11 info-de@origene.com

SM2096HRP

Monoclonal Antibody to Rat Lambda light chain - HRP

Quantity:	0.25 mg
Concentration:	1.0 mg/ml
Background:	All five immunoglobulin classes share the same basic four polypeptide chain structure of two heavy-chains and two light chains. There are five heavy chain types, and two light-chain types (Kappa and Lambda) both having a molecular weight of 22.5kDa. Any heavy-chain type can associate with either light-chain type, but on any immunoglobulin molecule both light-chains are of the same type. Kappa and Lambda consist of a variable region and a constant region and can easily be differentiated by the antigenic properties of the constant region. The ratio of Kappa to Lambda is 70:30 , the vast majority of which is bound to heavy-chain in immunoglobulin. In normal individuals low levels of free light-chain arepresent in serum (kappa, 1.6-15.2 mg/L; Lambda, 0.4-4.2mg/L), with the occurrence of multiple myeloma or other B-cell malignancies these levels can be greatly elevated and can be found at high levels in the urine (Bence-Jones proteins).
Host / Isotype:	Mouse / IgG1
Clone:	MARL-15
Immunogen:	Rat lambda light chain. Spleen cells from immunised BALB/c mice were fused with cells of the rat PAI-O myeloma cell line.
Format:	State: Liquid purified IgG Purification: Affinity chromatography Buffer System: PBS, pH7.4 containing 0.01% Thiomersal and 50% Glycerol Label: HRP – Horseradish Peroxidase
Applications:	ELISA. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Specificity:	This antibody recognises immunoglobulin lambda light chain. This antibody shows no cross-reactivity to Ig heavy chains or to kappa light chain. Species: Rat. Other species not tested.
Storage:	Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing. Shelf life: one year from despatch.

For research and in vitro use only. Not for diagnostic or therapeutic work. Material Safety Datasheets are available at www.acris-antibodies.com or on request.

OG/20121030 1/1