

SM24P**Armenian Hamster IgG Isotype Control - Purified**

Quantity:	0.25 mg
Concentration:	0.5 mg/ml
Host / Isotype:	Hamster / IgG
Clone:	HTK888
Immunogen:	Trinitrophenol - KLH
Format:	State: Liquid purified IgG fraction Purification: Ion Exchange Chromatography Buffer System: PBS, pH 7.2 Preservatives: 0.09% Sodium Azide
Applications:	Flow Cytometry. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Specificity:	This Hamster IgG Negative Control antibody is suitable for use as a Negative Control for the measurement of non-specific binding of Armenian Hamster monoclonal antibodies, of isotype IgG to Human, Mouse and Rat.
Storage:	Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing. Shelf life: one year from despatch.
General Readings:	1. Suzuki-Inoue K, Inoue O, Ding G, Nishimura S, Hokamura K, Eto K, et al. Essential in vivo roles of the C-type lectin receptor CLEC-2: embryonic/neonatal lethality of CLEC-2-deficient mice by blood/lymphatic misconnections and impaired thrombus formation of CLEC-2-deficient platelets. J Biol Chem. 2010 Aug 6;285(32):24494-507. doi: 10.1074/jbc.M110.130575. Epub 2010 Jun 4. PubMed PMID: 20525685.