

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850 UNITED STATES Phone: +1-888-267-4436 Fax: +1-301-340-8606 techsupport@origene.com

OriGene Technologies GmbH

Schillerstr. 5 32052 Herford GERMANY Phone: +49-5221-34606-0 Fax: +49-5221-34606-11 info-de@origene.com

SP1024F

Polyclonal Antibody to Human IgG/IgA/IgM FITC

Alternate names:	Human IgA, Human IgG, Human IgM
Quantity:	2 mg
Concentration:	1.0 mg/ml (after reconstitution)
Host:	Goat
Immunogen:	Polyclonal Human Ig.
Format:	 State: Lyophilized purified IgG fraction from Whole Serum Purification: Affinity Chromatography Buffer System: PBS, pH 7.2 Preservatives: 0.01% Sodium Azide Stabilizers: 1% BSA Label: FITC – Fluorescein Isothiocyanate Isomer 1 Reconstitution: Restore with 2.0 ml distilled water. Care should be taken during reconstitution as the protein may appear as a film at the bottom of the vial. We recommend that the vial is gently mixed after reconstitution.
Applications:	Flow Cytometry: Use 10 μ l of 0.1-0.5 μ g/ml diluted antibody to label 10 ⁶ cells in 100 μ l. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Specificity:	This antibody recognizes Human Ig, binding to all major classes (IgG, IgA, IgM). Species: Human. Other species not tested.
Storage:	Prior to reconstitution store at 2-8°C. Following reconstitution store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. This product is photosensitive and should be protected from light. Avoid repeated freezing and thawing. Shelf life: one year from despatch.

For research and in vitro use only. Not for diagnostic or therapeutic work. Material Safety Datasheets are available at www.acris-antibodies.com or on request.