

Polyclonal Antibody to DR3 (CT) Wsl-1, Apo-3, TRAMP, LARD

- Alternate names:** APO3, Apo-3, Apoptosis-inducing receptor AIR, Apoptosis-mediating receptor DR3, Apoptosis-mediating receptor TRAMP, DDR-3, DDR3, DR-3, DR3, LARD, Lymphocyte-associated receptor of death, TNFRSF12, TNFRSF25, WSL, WSL protein, WSL-1 protein, WSL1
- Catalog No.:** SP6253P
- Quantity:** 50 µg
- Concentration:** 0.5 mg/ml
- Host:** Rabbit
- Immunogen:** Rabbit anti-DR3 (CT) polyclonal antibody was raised against a peptide corresponding to amino acids 398 to 417 of human DR31,2. Peptide available as SP6253CP.
- Applications:** Western blot: 1/500 - 1/1000. Jurkat total cell lysate can be used as positive control and a 59 kDa band should be detected. Other applications not tested. Optimal dilutions of this antibody are dependent on conditions and should be determined by the user. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
- Specificity:** This antibody recognises a 59kD apoptosis related protein designated DR3 (death receptor-3). Apoptosis, or programmed cell death, occurs during normal cellular differentiation and development of multicellular organisms. Apoptosis is induced by certain cytokines including TNF and Fas ligand of the TNF family through their death domain containing receptors, TNFR1 and Fas. A novel cell death receptor was recently identified by several groups independently and designated DR3, Wsl-1, Apo-3, TRAMP and LARD1-5. The ligand for this novel cell death receptor has not yet been defined. DR3 is highly expressed in the tissues enriched in lymphocytes including PBL, thymus and spleen. Like TNFR1, DR3 induces apoptosis and NF- κ B activation. The antibody is human, mouse, and rat reactive and has no cross reaction to other death receptors.
- Storage:** Store the antibody at 4-8°C for one month or at -20°C for longer. Avoid repeated freezing and thawing. Shelf life: one year from despatch.
- General Readings:**
1. Chinnaiyan AM; O'Rourke K; Yu GL; Lyons RH; Garg M; Duan DR; Xing L; Gentz R; Ni J; Dixit VM. *Science*, 1996;274:990-2.
 2. Kitson J; Raven T; Jiang YP; Goeddel DV; Giles KM; Pun KT; Grinham CJ; Brown R; Farrow SN. *Nature*, 1996;384:372-5.
 3. Marsters SA; Sheridan JP; Donahue CJ; Pitti RM; Gray CL; Goddard AD; Bauer KD; Ashkenazi A. *Curr Biol*, 1996;6:1669-76.
 4. Bodmer JL; Burns K; Schneider P; Hofmann K; Steiner V; Thome M; Bornand T; Hahne M; Schroter M; Becker K; et al. *Immunity*, 1997;6:79-88.

For research and in vitro use only. Not for diagnostic or therapeutic work.

Material Safety Datasheets are available at www.acris-antibodies.com or on request.

Antibody Hotline - Technical Questions - Antibody Location Service
Free Call: 0800-2274746 (Germany only) - www.acris-antibodies.com



5. Screatton GR; Xu XN; Olsen AL; Cowper AE; Tan

For research and in vitro use only. Not for diagnostic or therapeutic work.

Material Safety Datasheets are available at www.acris-antibodies.com or on request.

Antibody Hotline - Technical Questions - Antibody Location Service
Free Call: 0800-2274746 (Germany only) - www.acris-antibodies.com