

## Acris Antibodies, Inc.

6815 Flanders Drive, Suite 140 San Diego, CA 92121 **UNITED STATES** 

Phone: +1-858-888-7900 Fax: +1-858-888-7904 US-info@acris-antibodies.com

## **Acris Antibodies GmbH**

Schillerstr. 5 32052 Herford **GERMANY** 

SP6255P

Phone: +49-5221-34606-0 Fax: +49-5221-34606-11 info@acris-antibodies.com

## Polyclonal Antibody to DR3 (NT) (WSL-1, APO-3, TRAMP, LARD)

APO3, Apo-3, Apoptosis-inducing receptor AIR, Apoptosis-mediating receptor DR3, Alternate names:

Apoptosis-mediating receptor TRAMP, DDR-3, DDR-3, DR-3, DR-3, LARD, Lymphocyte-

associated receptor of death, TNFRSF12, TNFRSF25, WSL, WSL protein, WSL-1 protein, WSL1

**Catalog No.:** SP6255P **Quantity:** 50 µg

**Concentration:** 0.5 mg/ml **Host:** Rabbit

Immunogen: Rabbit anti-DR3 (NT) polyclonal antibody was raised against a peptide corresponding to

amino acids 25 to 40 of human DR3 precusor1, 2. Peptide available as SP6255CP.

**Applications:** Western blot: 1/500 - 1/1000. Jurkat whole cell lysate can be used as positive control and a

59 kDa band can be detected. Other applications not tested. Optimal dilutions of this

antibody are dependent on conditions and should be determined by the user.

Other applications not tested. Optimal dilutions are dependent on conditions and should

be determined by the user.

**Specificity:** This antibody recognises a 59kD apoptosis related protein designated DR3 (death

receptor-3). Cross reacts with mouse.

Apoptosis, or programmed cell death, occurs during normal cellular differentiation and development of multicellular organisms. Apoptosis is induced by certain cytokines including TNF and Fas ligand of the TNF family through their death domain containing receptors, TNFR1 and Fas. A novel cell death receptor was recently identified by several groups independently and designated DR3, Wsl-1, Apo-3, TRAMP and LARD1-5. The ligand for this novel death receptor has been defined as TWEAK, also termed Apo3L. DR3 is highly expressed in the tissues enriched in lymphocytes including PBL, thymus and spleen. Like

TNFR1, DR3 induces apoptosis and NF-?B activation.

Store the antibody at 4-8°C for one month or at -20°C for longer. This product should be Storage:

stored undiluted. Avoid repeated freezing and thawing. Shelf life: one year from despatch.

1. Chinnaiyan AM; O'Rourke K; Yu GL; Lyons RH; Garg M; Duan DR; Xing L; Gentz R; Ni J; **General Readings:** 

Dixit VM. Science, 1996;274:990-2.

2. Kitson J; Raven T; Jiang YP; Goeddel DV; Giles KM; Pun KT; Grinham CJ; Brown R; Farrow

SN. Nature, 1996;384:372-5.

3. Marsters SA; Sheridan JP; Donahue CJ; Pitti RM; Gray CL; Goddard AD; Bauer KD;

Ashkenazi A. Curr Biol, 1996;6:1669-76.

4. Bodmer JL; Burns K; Schneider P; Hofmann K; Steiner V; Thome M; Bornand T; Hahne M;

Schroter M; Becker K; et al. Immunity, 1997;6:79-88. 5. Screaton GR; Xu XN; Olsen AL; Cowper AE; Tan

For research and in vitro use only. Not for diagnostic or therapeutic work.

Material Safety Datasheets are available at www.acris-antibodies.com or on request.