

Polyclonal Antibody to mSmac/DIABLO (CT)

Catalog No.:	SP6278P
Quantity:	50 µg
Concentration:	0.5 mg/ml
Host:	Rabbit
Immunogen:	A synthetic peptide corresponding to amino acids 222 to 237 of murine Smac/DIABLO (2).

Format: This antibody is supplied as liquid immunoaffinity purified immunoglobulin fraction in Phosphate buffered saline with 0.02% sodium azide as preservative.

Applications:	Suitable for Western blot (0.5 - 1 µg/ml, mouse heart tissue lysate can be used as positive control and a 25 kDa band can be detected). Other applications not tested. Optimal dilutions of this antibody are dependent on conditions and should be determined by the user. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
----------------------	---

Specificity:	The inhibitor of apoptosis proteins (IAPs) regulate programmed cell death by inhibiting members of the caspase family of enzymes. A novel mammalian protein that binds to IAPs and neutralizes the inhibitory effect of IAPs on caspases was recently identified and designated Smac/DIABLO (1, 2). Smac/DIABLO is a mitochondrial protein that is released along with cytochrome c during apoptosis and activates cytochrome c/Apaf-1/caspase-9 pathway. Analysis of the structural basis of Smac/DIABLO reveals that the N-terminal amino acids are required for binding of Smac/DIABLO to IAPs and activation of caspases (3-6). Smac/DIABLO is expressed in a variety of human and mouse tissues (1, 2). This antibody detects mSmac/DIABLO (CT) in human, mouse and rat.
---------------------	---

Storage:	Store the antibody undiluted at 4-8°C for one month or at -20°C for longer. Avoid repeated freezing and thawing. Should this product contain a precipitate we recommend microcentrifugation before use. Shelf life: one year from despatch.
-----------------	---

General Readings:	<ol style="list-style-type: none"> 1. Du C, Fang M, Li Y, Li L, Wang X. Smac, a mitochondrial protein that promotes cytochrome c-dependent caspase activation by eliminating IAP inhibition. <i>Cell</i>. 2000; 102(1):33-42. 2. Verhagen AM, Ekert PG, Pakusch M, Silke J, Connolly LM, Reid GE, Moritz RL, Simpson RJ, Vaux DL. Identification of DIABLO, a mammalian protein that promotes apoptosis by binding to and antagonizing IAP proteins. <i>Cell</i>. 2000; 102(1):43-53. 3. Srinivasula SM, Datta P, Fan XJ, Fernandes-Alnemri T, Huang Z, Alnemri ES. Molecular Determinants of the Caspase-promoting Activity of Smac/DIABLO and Its Role in the Death Receptor Pathway. <i>J Biol Chem</i>. 2000; 275(46):36152-36157. 4. Chai J, Du C, Wu JW, Kyn S, Wang X, Shi Y. Structural and biochemical basis of apoptotic activation by Smac/DIABLO. <i>Nature</i>. 2000; 406(6798):855-62. 5. Liu Z, Sun C, Olejniczak ET, Meadows RP, Betz SF, Oost T, Herrmann J, Wu JC, Fesik SW.
--------------------------	---

For research and in vitro use only. Not for diagnostic or therapeutic work.

Material Safety Datasheets are available at www.acris-antibodies.com or on request.

Antibody Hotline - Technical Questions - Antibody Location Service
Free Call: 0800-2274746 (Germany only) - www.acris-antibodies.com

Structural basis for binding of Smac/DIABLO to the XIAP BIR3 domain. Nature. 2000; 408(6815):1004-8.

6. Wu G, Chai J, Suber TL, Wu JW, Du C, Wang X, Shi Y. Structural basis of IAP recognition by Smac/DIABLO. Nature. 2000; 408(6815):1008-12.

SP6278P/AV0206

For research and in vitro use only. Not for diagnostic or therapeutic work.

Material Safety Datasheets are available at www.acris-antibodies.com or on request.

Antibody Hotline - Technical Questions - Antibody Location Service
Free Call: 0800-2274746 (Germany only) - www.acris-antibodies.com