

## OriGene Technologies Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850 UNITED STATES Phone: +1-888-267-4436 Fax: +1-301-340-8606 techsupport@origene.com

## TA309127 OriGene EU

Acris Antibodies GmbH Schillerstr. 5 32052 Herford GERMANY Phone: +49-5221-34606-0 Fax: +49-5221-34606-11 info@acris-antibodies.com

## Polyclonal Antibody to GABRA3 (N-term) - Aff - Purified

Alternate names:	GABA A Receptor subunit alpha-3, GABRA-3, Gamma-aminobutyric acid receptor subunit alpha-3
Catalog No.:	TA309127
Quantity:	0.1 ml
Background:	<i>Gamma</i> -aminobutyric acid (GABA) is the primary inhibitory neurotransmitter in the central nervous system. There are two major classes of GABA receptors: the GABAA and the GABAB subtype of receptors. GABAA-Rs are important therapeutic targets for a range of sedative, anxiolytic, and hypnotic agents and are implicated in several diseases including epilepsy, anxiety, depression, and sub-stance abuse. The GABAA-R is a multimeric subunit complex. To date six Alpha's, four Beta's and four Gamma's, plus alternative splicing variants of some of these subunits, have been identified (Olsen and Tobin, 1990; Whiting et al., 1999; Ogris et al., 2004). Injection in oocytes or mammalian cell lines of cRNA coding for Alpha-and beta-subunits results in the expression of functional GABAA-Rs sensitive to GABA. However, coexpression of a Gamma-subunit is required for benzodiazepine modulation. The various effects of the benzodiazepines in brain may also be mediated via different Alpha-subunits of the receptor (McKernan et al., 2000; Mehta and Ticku, 1998; Ogris et al., 2003).
Uniprot ID:	<u>P20236</u>
NCBI:	<u>NP_058765.1</u>
GenelD:	24947
Host / Isotype:	Rabbit / IgG
Immunogen:	Fusion protein from N-terminal region of the Alpha-3 subunit of rat GABAA Receptor.
Format:	<b>State:</b> Liquid purified Ig fraction. <b>Purification:</b> Affinity Chromatography. <b>Buffer System:</b> 10 mM HEPES (pH 7.5), 150 mM NaCl, 100 μg/ml BSA and 50% Glycerol.
Applications:	Western Blot: 1/1000. Immunohistochemistry: 1/100. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Specificity:	Specific for the ~51k Alpha-3 subunit of the GABAA receptor in Western blots. Labeling is absent in Alpha-3 subunit knockout animals.
Species Reactivity:	<b>Tested:</b> Rat and Mouse. <b>Expected from sequence similarity:</b> Human, Bovine, Canine, Zebrafish and non-Human Primates.

For research and in vitro use only. Not for diagnostic or therapeutic work. Material Safety Datasheets are available at www.acris-antibodies.com or on request.

OG/20160219

Acris Antibodies is now part of the OriGene family. Learn more at www.origene.com

	TA309127: Polyclonal Antibody to GABRA3 (N-term) - Aff - Purified
Storage:	Store the antibody undiluted (in aliquots) at -20°C. Avoid repeated freezing and thawing. Shelf life: one year from despatch.
General Readings:	<ol> <li>McKernan RM, Rosahl TW, Reynolds DS, Sur C, Wafford KA, Atack JR, et al. Sedative but not anxiolytic properties of benzodiazepines are mediated by the GABA(A) receptor alpha1 subtype. Nat Neurosci. 2000 Jun;3(6):587-92. PubMed PMID: 10816315.</li> <li>Mehta AK, Ticku MK (1998) Prevalence of the GABAA receptor assemblies containing Alpha-1 subunit in the rat cerebellum and cerebral cortex as determined by immunoprecipitation: Lack of modulation by chronic ethanol administration. Mol Brain Res 67:194-199.</li> <li>Ogris W, Pöltl A, Hauer B, Ernst M, Oberto A, Wulff P, et al. Affinity of various benzodiazepine site ligands in mice with a point mutation in the GABA(A) receptor gamma2 subunit. Biochem Pharmacol. 2004 Oct 15;68(8):1621-9. PubMed PMID: 15451405.</li> <li>Olsen RW, Tobin AJ (1990) Molecular biology of GABAA receptors. FASEB 4:1469-1480.</li> <li>Pöltl A, Hauer B, Fuchs K, Tretter V, Sieghart W. Subunit composition and quantitative importance of GABA(A) receptor subtypes in the cerebellum of mouse and rat. J Neurochem. 2003 Dec;87(6):1444-55. PubMed PMID: 14713300.</li> <li>Whiting PJ, Bonnert TP, McKernan RM, Farrar S, Le Bourdellès B, Heavens RP, et al. Molecular and functional diversity of the expanding GABA-A receptor gene family. Ann N Y Acad Sci. 1999 Apr 30;868:645-53. PubMed PMID: 10414349.</li> </ol>
Pictures:	Western blot of mouse brain lysates from wild type (Control) and $\hat{l} \pm \cdot \text{knockout 3}$ $(\hat{l} \pm \text{K}/\text{O})$ animals showing specific immunolabeling of the ~51k $\hat{l} \pm \cdot \text{subunit}$ of the 3 GABA -R. The labeling was absent from a lysate prepared from $\hat{l} \pm \cdot \text{knockout A3}$ animals. GABA $\alpha_3 - 43$

4



OG/20160219