

## Rabbit Polyclonal anti-TFII-I

<b>Catalog No.:</b>	VN55A-05
<b>Quantity:</b>	0.5 mg
<b>Background:</b>	This gene encodes a multifunctional phosphoprotein with roles in transcription and signal transduction. It is deleted in Williams-Beuren syndrome, a multisystem developmental disorder caused by the deletion of contiguous genes at chromosome 7q11.23. Four transcript variants encoding different isoforms have been found for this gene.
<b>Host / Isotype:</b>	Mouse / IgM
<b>Clone:</b>	N55-A
<b>Immunogen:</b>	The epitope recognized by this antibody maps to a region between residue 985 and 998 of human general transcription factor II, i using the numbering given in entry NP_127492.1 (GeneID 2969).
<b>Format:</b>	<b>Purification:</b> Immunogen affinity purified <b>Buffer System:</b> 0.09% Sodium Azide
<b>Applications:</b>	Immunohistochemistry-Paraffin, This antibody is useful for immunohistochemistry on paraffin sections. Epitope exposure with citrate buffer exposure is recommended. Epitope exposure with citrate buffer will enhance staining. Likely to work with frozen sections. In some cases, the antibody may be diluted further than indicated., Immunohistochemistry-Paraffin 1:100-1:500 Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
<b>Specificity:</b>	, TFII-I <b>Species:</b> Human, Mouse Others not tested.
<b>Storage:</b>	Store at 4C. Do not freeze.
<b>General References:</b>	

**For research and in vitro use only. Not for diagnostic or therapeutic work.**

Material Safety Datasheets are available at [www.acris-antibodies.com](http://www.acris-antibodies.com) or on request.

Antibody Hotline - Technical Questions - Antibody Location Service  
Free Call: 0800-2274746 (Germany only) - [www.acris-antibodies.com](http://www.acris-antibodies.com)