



VMRD, Inc.

PO Box 502, Pullman, WA 99163 USA

Telephone: + 1 (509) 334-5815

Fax: + 1 (509) 332-5356



E-mail: vmrd@vmrd.com

Web site: <http://www.vmrd.com>

Certificate of Analysis

Monoclonal Antibody

Cell Line:	TH97A
Specificity:	CD1b
Isotype:	IgG _{2a}
% Labeled Thymocytes:	70
Species Reactivity:	Cattle, goat, sheep, Cape buffalo, white tail deer, mule deer. Does not react with swine, cat, dog, ferret, mink, llama or camel.
Known Applications:	Flow cytometry, immunohistochemistry, cell function.



Profile
obtained
with bovine
leukocytes.

Description and Handling: This monoclonal antibody is produced as mouse ascites fluid, clarified by centrifugation, and filtered through a 0.2 µM filter. The concentration is 1.0 mg/ml in phosphate-buffered saline, preserved with sodium azide.

Storage: When the vial is stored at 2-8°C, it should be stable for one year.

FOR IN VITRO LABORATORY USE ONLY.

WARRANTY: VMRD, Inc. warrants that this product is as described in the quantity and contents stated on the label at the time of delivery to the customer. NO OTHER WARRANTIES, EXPRESS OR IMPLIED, ARE MADE BEYOND THE LABEL DESCRIPTION, INCLUDING WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR USE. Remedy is limited to replacement of the product or refund of the purchase price. VMRD, Inc. is not liable for property damage, personal injury, or economic loss caused by the product. The information listed in this information sheet is provided for reference only, and should not be substituted for the user's own incoming material quality control.

References:

- Sager, H., *et al.* Macrophage-parasite relationship in theileriosis. Reversible phenotypic and functional dedifferentiation of macrophages infected with *Theileria annulata*. *J. Leukocyte Biol.* 61(4):459-468 (Apr. 1997).
- Naessens, J., and J. Hopkins. Introduction and summary of workshop findings. *Vet. Immunol. Immunopathol.* 52:213-235 (1996).
- Davis, W.C., and R.A. Larsen. Genetic markers identified by immunogenetic methods. *Am. Fish. Soc. Symp.* 7:521-540 (1989).
- MacHugh, N.D., *et al.* Characterization of a bovine thymic differentiation antigen analogous to CD1 in the human. *Scand. J. Immunol.* 27(5):541-547 (May 1988).
- Davis, W.C., *et al.* The development and analysis of species specific and cross reactive monoclonal antibodies to leukocyte differentiation antigens and antigens of the major histocompatibility complex for use in the study of the immune system in cattle and other species. *Vet. Immunol. Immunopathol.* 15(4):337-376 (July 1987).